

Albac® contains feed grade zinc bacitracin equivalent to 150 grams (15%) bacitracin per kilogram.

INDICATIONS FOR USE

- 1. Supporting the prevention and control of necrotic enteritis caused or complicated by Clostridium spp.
- 2. Increased weight gain and improved feed efficiency.
- 3. Increased egg production and improved feed efficiency in laying hens.

KEY POINTS

- Albac® helps lowering intestinal counts of *C. perfringens* as well as other pathogenic bacteria, this promotes a healthy gut flora and higher nutrient absorption.
- The only bacitracin available for use in all ages of laying hens.
- Not absorbed from the intestinal tract, thus no withdrawal period is required prior to slaughter.
- Consistent performance after long time use.
- Effective in extreme weather conditions.

USERS INFORMATION: ALBAC® 15%

Active Drug Ingredient		Each kilogram contains feed grade zinc bacitracin equivalent to 150 g bacitracin (Master Standard).	
Composition		The zinc salt of a dried fermentation product obtained by culturing <i>Bacillus licheniformis</i> Tracy on media adapted for microbiological production of bacitracin and calcium carbonate.	
For Use in Manufactured Medicated Feeds			
Species	Grams bacitracin per tonne of Feed	Albac® dosage	Indications for Use
Broiler	5 - 50	dose between 35 - 335 gram /mton	For increased weight gain and improved feed efficiency
chickens	50	dose of 335 gram/mton	Supporting the prevention of necrotic enteritis. Medicated feed should be fed to poultry during the period when they are at most risk of the disease, which is usually between 2 and 12 weeks
	200	dose of 1335 gram/mton	Supporting the treatment of necrotic enteritis: add Albac® for 5 – 7 days
Laying hens	15 - 100	dose between 100-667 gram/mton	For increased weight gain and improved feed efficiency
Growing- finishing swine	10-50	dose between 66-335 gram/mton	For increased weight gain and improved feed efficiency Supports the prevention of enteritic disorders caused by <i>Clostridiaceae</i>

Can be used in combination with our registered feed additives and coccidiostats.*

HANDLING

- Albac® is packaged in a 25 kg, multilayer paper bag with protective barrier ply.
- Store in a cool (<25°C), dry place and keep package closed to avoid contamination.
- Expiry data is 24 months from the date of manufacture.
- For use in dry feed only.
- Not for use in feeds for breeder chickens.

*Specific claims vary by country.

Please contact your Huvepharma representative or distributor for more information.

References available on request.





